





















- Thompson-Brenner, H., Shingleton, R. M., Thompson, D. R., Satir, D. A., Richards, L. K., Pratt, E. M., & Barlow, D. H. (2016). Focused vs. broad enhanced cognitive behavioral therapy for bulimia nervosa with comorbid borderline personality: A randomized controlled trial. *International Journal of Eating Disorders*, 49(1), 36-49.
- Treasure, J., Sepulveda, A. R., MacDonald, P., Whitaker, W., Lopez C., Zabala, M., Kyriacou, O., & Todd, G. (2008). The assessment of the family of people with eating disorders. *European eating disorder review*, 16(4), 247-255.
- Treasure, J., Stein, D., & Maguire, S. (2015). Has the time come for a staging model to map the course of eating disorders from high risk to severe enduring illness? An examination of the evidence. *Early intervention in psychiatry*, 9(3), 173-184.
- Trompeter, N., Bussey, K., Forbes, M.K. & Mitchison, D. (2021). Emotion dysregulation within the CBT-E model of eating disorders: a narrative review. *Cognitive Therapy and Research*, 45(6), 1021-1036.
- Vitousek, K. B., & Hollon, S. D. (1990). The investigation of schematic content and processing in eating disorders. *Cognitive therapy and research*, 14(2), 191-214.
- Wells, A. (2008). *Metacognitive therapy for anxiety and depression*. The Guilford Press, New York.
- Westwood, H., Kerr-Gaffney, J., Stahl, D., & Tchanturia, K. (2017). Alexithymia in eating disorders: Systematic review and meta-analyses of studies using the Toronto Alexithymia Scale. *Journal of psychosomatic research*, 99, 66-81.
- Wilberg, T., Pedersen, G., Bremer, K., Johansen, M. S., & Kvarstein, E. H. (2023). Combined group and individual therapy for patients with avoidant personality disorder: A pilot study. *Frontiers in Psychiatry*, 14, 1181686.
- Wilson, G.T., Fairburn, C.C., Agras, W.S., Walsh, B.T. & Kraemer, H. (2002). Cognitive-behavioral therapy for bulimia nervosa: time course and mechanisms of change. *Journal of Consulting and Clinical Psychology*, 70(2), 267-274.
- Wonderlich, S. A., Peterson, C. B., Crosby, R. D., Smith, T. L., Klein, M. H., Mitchell, J. E., & Crow, S. J. (2014). A randomized controlled comparison of integrative cognitive-affective therapy (ICAT) and enhanced cognitive-behavioral therapy (CBT-E) for bulimia nervosa. *Psychological medicine*, 44(3), 543-553.
- Zipfel, S., Wild, B., Groß, G., Friederich, H. C., Teufel, M., Schellberg, D., & ANTOP Study Group. (2014). Focal psychodynamic therapy, cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. *The Lancet*, 383(9912), 127-137.
- Zipfel, S., Giel, K. E., Bulik, C. M., Hay, P., & Schmidt, U. (2015). Anorexia nervosa: aetiology, assessment, and treatment. *The lancet psychiatry*, 2(12), 1099-1111.

Non-commercial use only