

































- tween the patient's and therapist's view. *European Eating Disorders Review*, 15(5), 357–365. doi: 10.1002/erv.768
- Walfish, S., McAlister, B., O'Donnell, P., & Lambert, M. J. (2012). An investigation of self-assessment bias in mental health providers. *Psychological Reports*, 110(2), 639–644. doi: 10.2466/02.07.17.PR0.110.2.639-644
- Wampold, B. E. (2015). How important are the common factors in psychotherapy? An update. *World Psychiatry*, 14(3), 270–277. doi: 10.1002/wps.20238
- Wampold, B. E., & Owen, J. (2021). Therapist effects: History, methods, magnitude, and characteristics of effective therapists. In *Bergin and Garfield's handbook of psychotherapy and behavior change: 50th anniversary edition, 7th ed.* (pp. 297–326). John Wiley & Sons, Inc.
- Warren, C. S., Schafer, K. J., Crowley, M. E. J., & Olivardia, R. (2013). Treatment Providers With a Personal History of Eating Pathology: A Qualitative Examination of Common Experiences. *Eating Disorders*, 21(4), 295–309. doi: 10.1080/10640266.2013.797318
- Wasil, A., Venturo-Conerly, K., Shingleton, R., & Weisz, J. (2019). The motivating role of recovery self-disclosures from therapists and peers in eating disorder recovery: Perspectives of recovered women. *Psychotherapy*, 56(2), 170–180. doi: 10.1037/pst0000214
- Wells, G. A., Brodsky, L., O'Connell, D., Shea, B., Henry, D., Mayank, S., & Tugwell, P. (2003, October). An evaluation of the Newcastle Ottawa Scale: an assessment tool for evaluating the quality of non-randomized studies. In *XI International Cochrane colloquium book of abstracts* (p. 26). Spain: Barcelona.
- Werbart, A., Gråke, E., & Klingborg, F. (2023). Deadlock in psychotherapy: A phenomenological study of eight psychodynamic therapists' experiences. In N. Gazzola, S. Iwakabe, & S. Knox, *Therapeutic Failures in Psychotherapy* (1a ed., pp. 28–46). Routledge. doi: 10.4324/9781003439219-3
- Werz, J., Voderholzer, U., & Tuschen-Caffier, B. (2022). Alliance matters: But how much? A systematic review on therapeutic alliance and outcome in patients with anorexia nervosa and bulimia nervosa. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 27(4), 1279–1295. doi: 10.1007/s40519-021-01281-7
- Whitney, J., Currin, L., Murray, J., & Treasure, J. (2012). Family Work in Anorexia Nervosa: A Qualitative Study of Carers' Experiences of Two Methods of Family Intervention. *European Eating Disorders Review*, 20(2), 132–141. doi: 10.1002/erv.1077
- Wilfley, D. E., Welch, R. R., Stein, R. I., Spurrell, E. B., Cohen, L. R., Saelens, B. E., Douchis, J. Z., Frank, M. A., Wiseman, C. V., & Matt, G. E. (2002). A Randomized Comparison of Group Cognitive-Behavioral Therapy and Group Interpersonal Psychotherapy for the Treatment of Overweight Individuals With Binge-Eating Disorder. *Archives of General Psychiatry*, 59(8), 713. doi: 10.1001/archpsyc.59.8.713
- World Health Organization. (1992). *The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines (Volume 1)*. World Health Organization.
- Wright, K. M., & Hacking, S. (2012). An angel on my shoulder: A study of relationships between women with anorexia and healthcare professionals: An angel on my shoulder. *Journal of Psychiatric and Mental Health Nursing*, 19(2), 107–115. doi: 10.1111/j.1365-2850.2011.01760.x
- Zaitsoff, S., Pullmer, R., Menna, R., & Geller, J. (2016). A qualitative analysis of aspects of treatment that adolescents with anorexia identify as helpful. *Psychiatry Research*, 238, 251–256. doi: 10.1016/j.psychres.2016.02.045
- Zaitsoff, S., Yiu, A., Pullmer, R., Geller, J., & Menna, R. (2015). Therapeutic engagement: Perspectives from adolescents with eating disorders. *Psychiatry Research*, 230(2), 597–603. doi: 10.1016/j.psychres.2015.10.010

Non-commercial use only

Online supplementary material:

Supplementary Table 1. Quality assessment of included qualitative studies.

Supplementary Table 2. Quality assessment of included cohort studies and mixed methods.